



Tarım Bilimleri Dergisi
Tar. Bil. Der.

Dergi web sayfası:
www.agri.ankara.edu.tr/dergi

Journal of Agricultural Sciences

Journal homepage:
www.agri.ankara.edu.tr/journal

Effects of Different Quality Roughage Supply on Performance of Holstein Calves during Preweaning Period

Ismail ÜLGER^a, Mahmut KALİBER^a, Selma BÜYÜKKILIÇ BEYZİ^a, Yusuf KONCA^a

^aErciyes University, Faculty of Agriculture, Department of Animal Science, Kayseri, TURKEY

ARTICLE INFO

Research Article

Corresponding Author: Mahmut KALİBER, E-mail: mkaliber@gmail.com, Tel: +90 (507) 540 11 45

Received: 18 January 2016, Received in Revised Form: 29 February 2016, Accepted: 08 March 2016

ABSTRACT

The present study was conducted to investigate the effects of two different quality roughage (wheat straw and alfalfa hay) supplies on live weight gain, growth, feed intake and feed efficiency of Holstein calves during the preweaning period. With this purpose, 30 male and female Holstein calves were divided into two treatment groups (15 calves in each group) based on gender and birth weights. The first group was fed with a total mixed diet containing 80% calf starter and 20% alfalfa hay (chopped in 1-2 cm pieces) as a high quality roughage source and the second group was fed with a total mixed diet containing 80% calf starter and 20% wheat straw (chopped in 1-2 cm pieces) as a low quality roughage source during the preweaning period. The results revealed that feeding with high quality roughage (alfalfa hay) decreased feed intake ($P<0.05$) but didn't affect live weight and growth parameters ($P>0.05$) at the moment of weaning after 8 week preweaning period. Because of lower feed consumption without impairing calf performance and improved feed conversation efficiency, it can be suggested that high quality roughage could be supplied to calves in addition to calf starter feed during the preweaning period but meanwhile economic parameters like feed costs must be taken into account.

Keywords: Holstein; Calf; Weaning; Alfalfa; Wheat straw; Growth performance

Sütten Kesim Öncesi Dönemde Farklı Kalitedeki Kaba Yemlerle Beslemenin Holstein Buzağuların Performansı Üzerine Etkileri

ESER BİLGİSİ

Araştırma Makalesi

Sorumlu Yazar: Mahmut KALİBER, E-posta: mkaliber@gmail.com, Tel: +90 (507) 540 11 45

Geliş Tarihi: 18 Ocak 2016, Düzeltmelerin Gelişi: 29 Şubat 2016, Kabul: 08 Mart 2016

ÖZET

Bu çalışmada, Holstein buzağularında süttten kesim öncesi dönemde iki farklı kalitedeki kaba yem kullanımının (kıyılmış halde kuru yonca otu ve buğday sapı) büyüme, gelişme, yem tüketimi ve yemden yararlanma özellikleri üzerindeki etkileri araştırılmıştır. Bu amaçla toplam 30 baş erkek ve dişi Holstein buzağı cinsiyet ve doğum ağırlıkları göz önünde bulundurularak her birinde 15 baş olmak üzere 2 deneme grubuna dağıtılmıştır. Birinci muamele grubunun % 80

oranında buzağı başlangıç yemi ve kaliteli kaba yem kaynağı olarak % 20 oranında 1-2 cm uzunluğundaki kıyılmış yonca kuru otu, ikinci muamele grubunun ise % 80 oranında buzağı başlangıç yemi ve düşük kaliteli kaba yem kaynağı olarak % 20 oranında 1-2 cm uzunluğundaki kuru buğday sapı kullanılarak hazırlanan toplam yem karışımını (TMR) 8 haftalık süt ile besleme periyodu boyunca tüketmeleri sağlanmıştır. Çalışma sonucunda, buzağuların süttan kesim öncesi dönemde buzağı başlangıç yeminin yanında kaliteli kaba yemler ile beslenmesinin yem tüketimini azalttığı ($P<0.05$) ancak buna rağmen 8 haftalık süt ile besleme programı sonunda süttan kesim ağırlıklarını ve iskelet gelişimlerini etkilemediği ($P>0.05$) tespit edilmiştir. Bu noktadan hareketle, buzağuların performanslarının etkilenmeden daha az yem ile süttan kesilmelerine olanak sağlaması ayrıca yem çevirim kabiliyetlerini artırması ($P<0.05$) gibi faydaları göz önünde bulundurularak süttan kesim öncesi dönemde buzağuların kesif yeme ek olarak kaliteli kaba yemler ile beslenmesinin ekonomik faktörlerin de göz önünde bulundurulması koşuluyla uygun olacağı sonucuna varılmıştır.

Anahtar Kelimeler: Siyah Alaca; Buzağı; Süttan kesme; Yonca; Saman; Performans

© Ankara Üniversitesi Ziraat Fakültesi

1. Introduction

In dairy operations, proper growth and feeding of newborn calves are significant issues for replacing the dairy cows with the young ones. While milk constitutes about 60% of total facility income, calf and stock capital gain constitute 40% of total income. In other words, number of calves raised in the facility has a great impact on profitability of the operation (Erez 2011).

Including calves, proventriculus (rumen, reticulum and omasum) of young ruminants is not developed sufficiently during the initial periods of their lives. Early weaning without much influencing the calf performance may increase marketable milk production of the operation and protein supplements to the feeds cost less than the protein supplied with the milk (Turgut et al 1998). Thus, roughage and concentrate feeds should be used efficiently and proventriculus of calves should be developed as fast as possible to stimulate rumen development. Proventriculus development will be possible with the use of solid feeds (Göncü et al 2010). Several researchers indicated that roughages could be used for normal development of rumen throughout the preweaning period (Beharka et al 1998; Coverdale et al 2004; Waterman 2005; Van Ackeren et al 2009). Usage of roughages during the preweaning period may have stimulating impact on rumination (Lyford & Huber 1988), improve rumen action, volume and muscularity (Tamate et al 1962) and support epithelium and papilla development

(Wardrop 1961). Parallel to this, some researchers have recommended the addition of roughage to the starter for optimum rumen development (Anderson et al 1987; Greenwood et al 1997), but some other researchers (Heinrichs & Lesmeister 2000; Lizieire et al 2002; Harris & Shearer 2003; Jones & Heinrichs 2007) did not recommend roughage during preweaning period. However, calf rearing without roughage was seen possible only for short-term weaning programs (<30 days) (Hibbs et al 1956; Leibholz 1975; Porter et al 2007).

It was indicated in studies investigating the effects of roughage forms of the ration that roughage forms did not have any significant impacts on growth performance of the calves (Leaver 1973; Kang & Leibholz 1973). It was also indicated in a previous study that wheat straw chopped at different lengths (1.5, 4.8 and 7.9 mm) did not yield significant differences in growth performance of calves (Pickard et al 1969).

Grass and legume hay and straw are common sources of fiber in the diet of the young calf. The quality of these forage sources greatly affects the palatability and the digestibility of the diet (Booth 2003). Differences in feed quality attributes were numerically assessed thorough relative feed value. Previous researchers investigated relative feed values of dry alfalfa hay and wheat straw (Yavuz 2005) and assessed digestibility and energy levels of different roughages (Denek & Deniz 2004). Güngör et al (2008) reported higher crude nutrient contents

for alfalfa than for wheat straw and indicated alfalfa as a quality roughage source.

There are several studies carried out about the effects of different preweaning milk feeding periods (Ozkaya & Toker 2012; Kocyigit et al 2015; Ayaşan et al 2015), feeding with roughage and without roughage (Göncü et al 2010; Daneshvar et al 2015) and different feed forms (Montoro et al 2013; Jahani-Moghadam et al 2015) and the studies about the effects of different roughage sources on calf performance have come into prominence (Zanton & Heinrichs 2009; Castells et al 2012; Terré et al 2013). However, despite all these studies, effects of roughage quality on calf performance are not clear, yet. Therefore, the present study was conducted to investigate the effects of roughage quality in 8 weeks preweaning period on calf performance.

2. Material and Methods

The present study was conducted with 30 Holstein calves in a livestock facility located in Kayseri province of Turkey (latitude: 38.73695, longitude: 35.49683). Two treatment groups were formed with 15 calves consuming poor and high quality roughage. Calves were tried to be placed in treatment groups as much equally as possible based on gender (8 males and 7 females in each treatment group), age (all of the calves were born from the mothers at the same age within 24 hours), amount of colostrum intake and live weight (Table 1). Experiments were implemented throughout 8 weeks of preweaning period. The first treatment group (dry alfalfa hay-DAH) was supplied with 80% calf starter feed obtained from a commercial feed factory in Kayseri and 20% dry alfalfa hay chopped at 1-2 cm pieces as a quality roughage source in addition to milk from the 14th day of birth. The second group (wheat

straw-WS) was supplied with total mixed ration (TMR) composed of 80% calf starter feed and 20% dry wheat straw chopped at 1-2 cm pieces as a poor quality roughage source. All calves were fed *ad libitum*.

Calves were housed in individual pens throughout the experiments and straw bedding was used over pen floors. The calves stayed with their mothers in the first day. They took colostrum within maximum 2 hours from the birth and colostrum was supplied to calves for 3 days with a feeding bottle. From the 3rd day till weaning, the calves consumed a total of 4 liters whole milk (the average values obtained from the analyses performed on morning milk samples taken in every two weeks by using FOSS FT120 milk analysis device: fat 3.81%, protein 3.09%, lactose 4.49% and fat-free dry matter 8.19%) daily in two diets with a nipple bucket. All calves consumed a total of 212 liters milk throughout 8 weeks of milk-feeding period. Clean drinking water was supplied *ad libitum*.

Calves were weighed in every two weeks starting from the birth with an electronic scale (100 kg ± 0.001 kg) to calculate live weight gains. Feeds (hay and concentrated feed) were weighed every day to determine daily feed consumption of the calves. Skeletal growth was determined through measuring the chest girths, withers heights and hip widths in accordance with the methods specified in literature (Heinrichs & Lammers 1998). Feed conversion rates were calculated by dividing daily dry matter intake (g) with daily live weight gain (g). The changes in live weights were determined by subtracting birth weights from the live weights at the end of experiments (56th day).

Analyses of dry matter (DM), crude ash (CA), crude protein (CP) and crude fat (CF) contents of

Table 1- Birth weights and colostrum intake of calves

Parameters	Groups		SE	Significance
	WS	DAH		
Birth weight, kg	39.50	39.60	0.91	ns
Total Colostrum intake, L	16.30	16.80	0.95	ns

DAH, dry alfalfa hay; WS, wheat straw; SE, standard error; ns, not significant

roughage sources (wheat straw and alfalfa hay) were carried out in accordance with AOAC (1998); acid detergent fiber (ADF) and neutral detergent fiber (NDF) analyses were carried out in accordance with Van Soest et al (1991); metabolic energy (ME) and organic matter digestibility (OMD) values were calculated with the equation developed by Menke et al (1979) by using the amount of gases released after 24 hours fermentation through in vitro gas production technique. Rumen fluid was obtained from two fistulated sheep fed twice daily with a diet containing alfalfa hay (60%) and concentrate feed (40%). About 0.200 g dry samples were weighed in triplicate into the calibrated 100 mL glass syringes. The syringes were pre-warmed to 39 °C before the injection of 30 mL rumen fluid-buffer mixture into each syringe followed by incubation in a water bath at 39 °C. Readings of gas production were recorded before incubation (0) and 24 h after incubation.

The metabolic energy (ME) (Mcal kg⁻¹ DM) of feedstuffs was calculated according to Menke et al (1979) (Equation 1).

$$ME = (2.20 + 0.1136 \times GP + 0.0057 \times CP + 0.00029 \times CF) \times 4.184 \quad (1)$$

Where; GP, 24 h net gas production (mL 200 mg⁻¹); CP, crude protein (%) and CF, crude fat (%).

The organic matter digestibility (OMD) of feedstuffs was calculated according to Menke et al (1979) (Equation 2).

$$OMD (\%) = 14.88 + 0.889 \times GP + 0.45 \times CP + 0.0651 \times CA \quad (2)$$

Where; GP, 24 h net gas production (mL 200 mg⁻¹); CP, crude protein (%); CA, crude ash (%).

The equations specified by Van Dyke & Anderson (2000) were used to determine relative feed value (RFV) (Table 2) of roughage sources used in experiments. Relative feed values were calculated by using digestible dry matter (DDM) and dry matter intake (DMI) values. Digestible dry matter was calculated by using ADF and dry matter intake was calculated by using NDF values. Results are provided in Table 2.

Table 2- Nutritive values and quality attributes of calf starter feed, alfalfa hay and wheat straw

Nutrients	Feed materials		
	Calf starter feed*	Alfalfa	Wheat straw
Crude protein, %	18	18.06	3.74
Dry matter, %	88	91.72	90.14
Crude ash, %	8.00	9.78	8.02
Crude fat, %	3.60	2.52	1.49
ADF, %	18	36.09	52.37
NDF, %	30	44.41	80.14
ME, kcal kg ⁻¹	2800	1810	1140
OMD, %	-	58.81	32.87
DDM, %	-	60.79	48.10
DMI, %	-	2.70	1.50
RFV	-	127.20	55.92

*, values were received from the producer; ADF, acid detergent fiber; NDF, neutral detergent fiber; ME, metabolic energy; OMD, organic matter digestibility; DDM, digestible dry matter= 88.9-(0.779×ADF); DMI, dry matter intake= 120/NDF; RFV, relative feed value= DDM×DMI×0.775

Descriptive statistics for studied variables were presented as mean and standard errors. Experimental results were subjected to the independent samples t-test to determine whether or not the differences between the groups were significant. Significance level was considered as 5% and SPSS statistical software was used in all statistical analyses (SPSS 2002).

3. Results and Discussion

Daily feed consumption (DFC), initial live weights (ILW), mid-experiment (4th week) live weights (MLW), final live weights (FLW), daily live weight gains (LWG), feed conversion ratios (FCR), initial chest girths (ICG), mid-experiment (4th week) chest girths (MCG), final chest girths (FCG), initial hip widths (IHW), mid-experiment (4th week) hip widths (MHW), final hip widths (FHW), initial withers heights (IWH), mid-experiment (4th week) withers heights (MWH), final withers heights (FWH), changes in live weights (CLW), changes in chest girths (CCG), changes in hip widths (CHW) and changes in withers heights (CWH) of the calves fed with different quality roughages throughout 8 week preweaning period are provided in Table 3.

Table 3- Performance and growth parameters of calves

	<i>Groups</i>			<i>Significance</i>
	<i>WS</i>	<i>DAH</i>	<i>SE</i>	
<i>Live weight</i>				
ILW, kg	39.50	39.60	0.91	ns
MLW, kg	57.50	58.00	1.38	ns
FLW, kg	70.20	70.80	0.93	ns
LWG, g day ⁻¹	548.21	557.14	1.08	ns
CLW, kg	30.70	31.20	0.33	ns
<i>Feed consumption</i>				
DFC, g day ⁻¹	883.26	859.65	7.39	**
FCR, DFC LWG ⁻¹	1.61	1.54	0.03	*
<i>Chest girth</i>				
ICG, cm	78.60	79.80	1.53	ns
MCG, cm	89.50	91.85	1.36	ns
FCG, cm	98.00	100.80	1.97	ns
CCG, cm	19.40	21.00	1.08	ns
<i>Hip width</i>				
IHW, cm	19.15	19.50	0.73	ns
MHW, cm	21.85	22.40	0.93	ns
FHW, cm	26.50	27.40	0.94	ns
CHW, cm	7.35	7.90	0.67	ns
<i>Withers height</i>				
IWH, cm	75.50	76.00	1.23	ns
MWH, cm	78.70	80.00	1.24	ns
FWH, cm	84.20	86.45	1.87	ns
CWH, cm	8.70	10.45	1.25	ns

DAH, dry alfalfa hay; *WS*, wheat straw; *SE*, standard error; *, < 0.05; **, < 0.01; ns, not significant; ILW, initial live weight; MLW, mid-experiment (4th week) live weight; FLW, final live weight; LWG, live weight gain; CLW, changes in live weight; DFC, daily feed consumption; FCR, feed conversion ratio (contains only dry feed); ICG, initial chest girth; MCG, mid-experiment (4th week) chest girth; FCG, final chest girth; CCG, changes in chest girth; IHW, initial hip width; MHW, mid-experiment (4th week) hip width; FHW, final hip width; CHW, changes in hip width; IWH, initial withers height; MWH, mid-experiment (4th week) withers height; FWH, final withers height; CWH, changes in withers height

The differences in initial, mid-experiment and final live weights, daily live weight gains, changes in live weights, initial, mid-experiment and final chest girths, changes in chest girths, initial, mid-experiment and final hip widths, changes in hip widths, initial, mid-experiment and final withers heights and changes in withers heights of experimental groups were not found to be significant ($P>0.05$). It was reported in previous

studies (Greenwood et al 1997; Beharka et al 1998; Coverdale et al 2004; Suarez et al 2007; Castells et al 2012) that calf performance and live weights varied based on weaning age, type of roughage supply (*ad libitum* or TMR), form of roughage supply (hay or smaller pieces) and roughage (e.g., alfalfa hay, straw, grass hay, fresh grass, corn silage). In this study, similar growth performance and weaning weights of the calves fed with high

and low quality roughage were probably resulted from equal milk feeding (4 L day⁻¹) of treatment groups, relatively low ratios (20%) of roughage in TMR and relatively shorter treatment durations (14-56th days).

On the other hand, roughage quality had significant impacts on daily feed consumptions and feed conversion ratios ($P<0.05$). While the daily average feed consumption of the calves fed with alfalfa hay as a roughage was 859.65 g, the value was observed to be 883.26 g in treatment group fed with wheat straw ($P<0.05$). The calves fed with wheat straw throughout the preweaning period had higher feed consumptions than the calves fed with alfalfa hay. Complying with the findings of earlier studies (Beharka et al 1998; Booth 2003; Coverdale et al 2004; Suarez et al 2007), use of low quality roughage instead of high quality roughage in rations increased daily feed consumptions throughout the preweaning period. However, current findings conflict with those observed by Castells et al (2012) who observed greater DFC in calves fed with WS than DAH. However in that study, roughage was not supplied in TMR, they were supplied freely, thus the animals fed with WS consumed 5% roughage and the ones fed with DAH consumed 15% roughage. Therefore, the DGC values were higher than the present values. Since the wheat straw used in present study had 1.58% less dry matter content than alfalfa hay (Table 2), the difference in dry matter contents might have an increasing effect on feed consumption. Such a difference in feed consumptions might have also resulted from higher digestible nutrient content of alfalfa hay than wheat straw and consequently the calves fed with alfalfa hay was able to meet their nutritional needs with less feed consumption. Furthermore, increasing feed intakes with low quality roughage might be due to a bulky effect, which helps to stimulate rumination and improve rumen pH of young calves, instead of an increase of the NDF content of the diet (Booth 2003).

Feed conversion ratio of the treatment group fed with wheat straw (1.61) was higher than the

ratio in alfalfa consuming group (1.54) ($P<0.05$). Lower feed conversion ratios indicate better conversions, in other words less feed consumptions for unit live weight gain. Accordingly, while the daily feed consumption of the treatment group fed with alfalfa hay (DAH) was 859.65 g and the daily live weight gain of the group was 557.14 g, the values were respectively observed as 883.26 g (DFC) and 548.21 g (LWG) in treatment group fed with wheat straw (WS). Complying with the earlier findings (Leaver 1973; Greenwood et al 1997; Booth 2003; Coverdale et al 2004), feed conversion ability of the calves fed with high quality roughage throughout the preweaning period was better than the calves fed with low quality roughage. Again higher digestible nutrient and lower NDF content of alfalfa hay than wheat straw (Table 2) resulted in higher nutrient intake in alfalfa consumed calves than in wheat straw consumed calves, thus better feed conversions in alfalfa group.

4. Conclusions

Current findings revealed that calf feeding with quality roughage throughout the preweaning period in addition to concentrate feed decreased the feed consumption of the calves ($P<0.05$). However such a feeding program throughout the 8 weeks of milk-fed period did not significantly influence weaning weights and skeletal development of the calves ($P>0.05$). It was concluded that quality roughage supply in addition to concentrate feed was found to be a convenient approach to allow weaning of calves with less feed consumption without impairing their performances and to improve feed conversion abilities of the calves in an economical fashion ($P<0.05$). The present study can be considered as an initial step for further studies and as an indicator for producer-level practices. There is a need for further studies to investigate the rumen development of calves reared with different quality roughages, to assess further performance of calves and to evaluate the effects on dairy and beef cattle separately.

Abbreviations and Symbols			
<i>kg</i>	kilogram	<i>FLW</i>	Final live weight
<i>g</i>	gram	<i>FCG</i>	Final chest girth
<i>L</i>	liter	<i>FHW</i>	Final hip width
<i>mL</i>	milliliter	<i>FWH</i>	Final withers height
<i>m</i>	meter	<i>CLW</i>	Changes in live weight
<i>cm</i>	centimeter	<i>CCG</i>	Changes in chest girth
<i>mm</i>	millimeter	<i>CHW</i>	Changes in hip width
<i>DAH</i>	Dry alfalfa hay	<i>CWH</i>	Changes in withers height
<i>WS</i>	Wheat straw	<i>ADF</i>	Acid detergent fiber
<i>TMR</i>	Total mixed ration	<i>NDF</i>	Neutral detergent fiber
<i>ILW</i>	Initial live weight	<i>OMD</i>	Organic matter digestibility
<i>ICG</i>	Initial chest girth	<i>ME</i>	Metabolic energy
<i>IHW</i>	Initial hip width	<i>DDM</i>	Digestible dry matter
<i>IWH</i>	Initial withers height	<i>DMI</i>	Dry matter intake
<i>MLW</i>	Mid-experiment live weight	<i>RFV</i>	Relative feed value
<i>MCG</i>	Mid-experiment chest girth	<i>FCR</i>	Feed conversion ratio
<i>MHW</i>	Mid-experiment hip width	<i>DFC</i>	Daily feed consumption
<i>MWH</i>	Mid-experiment withers height		

References

- Anderson K L, Nagaraja T G, Morrill J L, Avery T B, Galitzer S J & Boyer J E (1987). Ruminant microbial development in conventionally or early-weaned calves. *Journal of Animal Science* **64**(4): 1215-1226
- AOAC (1998). Official methods of analysis. Association of Official Analytical Chemists, Arlington, Virginia, USA
- Ayaşan T, Hızlı H & Ünalın A (2015). Farklı sütten kesim yaşının Siyah Alaca buzağuların canlı ağırlık artışı, vücut ölçütleri ve süt maliyetine olan etkisi. *9. Ulusal Zooteknik Bilim Kongresi*: 3-5 Eylül, Konya, s. 650-658
- Beharka A A, Nagaraja T G, Morrill J L, Kennedy G A & Klemm R D (1998). Effects of form of the diet on anatomical, microbial and fermentative development of the rumen of neonatal calves. *Journal of Dairy Science* **81**: 1946-1955
- Booth J A (2003). Effect of forage addition to the diet on rumen development in calves. PhD Thesis, Iowa State University (Unpublished), USA
- Castells L, Bach A, Araujo G, Montoro C & Terré M (2012). Effect of different forage sources on performance and feeding behavior of Holstein calves. *Journal of Dairy Science* **95**(1): 286-293
- Coverdale J A, Tyler H D, Quigley J D & Brumm J A (2004). Effect of various levels of forage and form of diet on rumen development and growth in calves. *Journal of Dairy Science* **87**: 2554-2562
- Daneshvar D, Khorvash M, Ghasemi E, Mahdavi A H, Moshiri B, Mirzaei M, Pezeshki A & Ghaffari M H (2015). The effect of restricted milk feeding through conventional or step-down methods with or without forage provision in starter feed on performance of Holstein bull calves. *Journal of Animal Science* **93**: 3979-3989
- Denek N & Deniz S (2004). Ruminant beslemede yaygın olarak kullanılan kimi kaba yemlerin sindirilebilirlik ve metabolik enerji düzeylerinin in vitro yöntemlerle belirlenmesi. *Turkish Journal of Veterinary and Animal Sciences* **28**: 115-122
- Erez İ (2011). Siyah Alaca buzağularda erken sütten kesmenin performans üzerine etkileri. Yüksek lisans tezi, Çukurova Üniversitesi Fen Bilimleri Enstitüsü (Basılmamış), Adana
- Göncü S, Boğa M, Kılıç Ü, Görgülü M & Doran F (2010). Effects of feeding regime without roughage on performances and rumen development of calves during preweaning period. *Tarım Bilimleri Dergisi- Journal of Agricultural Sciences* **16**: 123-128

- Greenwood R H, Morrill J L, Titgemeyer E C & Kennedy G A (1997). A new method of measuring diet abrasion and its effect on the development of the forestomach. *Journal of Dairy Science* **80**: 2534-2541
- Güngör T, Başalan M & Aydoğan İ (2008). Kırıkkale yöresinde üretilen bazı kaba yemlerde besin madde miktarları ve metabolize olabilir enerji düzeylerinin belirlenmesi. *Ankara Üniversitesi Veteriner Fakültesi Dergisi* **55**: 111-115
- Harris B Jr & Shearer J K (2003). Feeding and management of young dairy calves. <http://edis.ifas.ufl.edu/ds117/> (Accessed: 24.10.2015)
- Heinrichs J & Lammers B (1998). Monitoring dairy heifer growth. PennState Publications, College of Agricultural Sciences, No: 5M498PS, USA
- Heinrichs A J & Lesmeister K (2000). Why you should hold off on feeding forage to calves. *Hoard's Dairyman* **145**: 638
- Hibbs J W, Conrad H R, Pouden W D & Frank N (1956). A high roughage system for raising calves based on early development of rumen function. VI. Influence of hay to grain ratio on calf performance, rumen development, and certain blood changes. *Journal of Dairy Science* **39**: 171-179
- Jahani-Moghadam M, Mahjoubi E, Yazdi M H, Cardoso F C & Drackley J K (2015). Effects of alfalfa hay and its physical form (chopped versus pelleted) on performance of Holstein calves. *Journal of Dairy Science* **98**(6): 4055-4061
- Jones C & Heinrichs J (2007). Early weaning strategies. http://www.extension.org/pages/early_weaning_strategies/ (Accessed: 20.10.2015)
- Kang H S & Leibholz J (1973). The roughage requirement of the early-weaned calf. *Animal Production* **16**: 195-203
- Kocuyigit R, Aydın R, Yanar M, Guler O, Diler A, Tuzemen N, Avci M, Ozyurek S & Kabakci D (2015). Effect of doses of direct-fed microbials plus exogenous fibrolytic enzymes supplementation on growth feed efficiency ratio and fecal consistency index of Brown Swiss and Holstein Friesian calves. *Indian Journal of Animal Research* **49**(1): 63-69
- Leaver J D (1973). Rearing dairy cattle. IV. Effect of concentrate supplementation on the live weight gain and feed intake of calves offered roughages ad libitum. *Animal Production* **17**: 43-52
- Leibholz J (1975). Ground roughage in the diet of the early weaned calves. *Animal Production* **20**: 93-100
- Lizieire R C, Cunha D N, Da F V, Martuscello J A & Campos O F (2002). Fornecimento de volumoso para bezerros pré-ruminantes. *Ciência Rural* **32**: 835-840
- Lyford S J & Huber J T (1988). Digestion, metabolism and nutrient needs in preruminants. In: DC Church (Ed), *The Ruminant Animal. Digestive Physiology and Nutrition*, Edition Prentice-Hall, Englewood Cliffs, NJ, USA, pp. 401
- Menke K H, Raab L, Salewski A, Steingass H, Fritz D & Schneider W (1979). The estimation of the digestibility and metabolizable energy content of ruminant feedstuffs from the gas production when they are incubated with rumen liquor. *The Journal of Agricultural Science* **93**: 217-222
- Montoro C, Miller-Cushon E K, DeVries T J & Bach A (2013). Effect of physical form of forage on performance, feeding behavior, and digestibility of Holstein calves. *Journal of Dairy Science* **96**: 1117-1124
- Ozkaya S & Toker M T (2012). Effect of amount of milk fed, weaning age and starter protein level on growth performance in Holstein calves. *Archiv Tierzucht* **55**: 234-244
- Pickard D W, Swan H & Lamming G E (1969). Studies on the nutrition of ruminants. IV. The use of ground straw of different particle sizes for cattle from twelve weeks of age. *Animal Production* **11**: 543-550
- Porter J C, Warner R G & Kertz A F (2007). Effects of fiber level and physical form of starter on growth and development of dairy calves fed no forage. *The Professional Animal Scientist* **23**: 395-400
- SPSS (2002). SPSS for windows, release 11.5.0. SPSS Inc, Chicago, Illinois, USA
- Suarez B J, Van Reenen C G, Stockhofe N, Dijkstra J & Gerrits W J J (2007). Effect of roughage source and roughage to concentrate ratio on animal performance and rumen development in veal calves. *Journal of Dairy Science* **90**: 2390-2403
- Tamate H, McGilliard A D, Jacobson N L & Getty R (1962). Effect of various dietaries on the anatomical development of the stomach in the calf. *Journal of Dairy Science* **45**: 408-420
- Terré M, Pedrals E, Dalmau A & Bach A (2013). What do preweaned and weaned calves need in the diet: A high fiber content or a forage source? *Journal of Dairy Science* **96**(8): 5217-5225
- Turgut L, Yanar M & Tüzemen N (1998). Kaba yem formunun Esmer buzağılarda büyüme ve yemden

- yararlanma özelliklerine etkileri. *Doğu Anadolu Tarım Kongresi*. Bildiriler; 14-18 Eylül, Erzurum, s. 835-845
- Van Ackeren C, Steingas H, Hartung K, Funk R & Drochner W (2009). Effect of roughage level in a total mixed ration on feed intake, ruminal fermentation patterns and chewing activity of early-weaned calves with ad libitum access to grass hay. *Animal Feed Science and Technology* **153**: 48-59
- Van Dyke N J & Anderson P M (2000). Interpreting a forage analysis. Alabama cooperative extension, Circular ANR-890, USA
- Van Soest P J, Robertson J B & Lewis B A (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* **74**: 3583-3597
- Wardrop I D (1961). Some preliminary observations on the histological development of the forestomachs of the lamb. 1: Histological changes due to age in the period from 46 days of foetal life to 77 days post-natal life. *The Journal of Agricultural Science* **57**: 335-341
- Waterman D F (2005). Sources of nutrients for milk replacers and dry starter feeds and what factors impact quality. NRAES-175, Cooperative Extension, pp. 96-115
- Yavuz M (2005). Bazı ruminant yemlerinin nispi yem değeri ve in vitro sindirim değerlerinin belirlenmesi. *Gaziosmanpaşa Üniversitesi Ziraat Fakültesi Dergisi* **22**(1): 97-101
- Zanton G I & Heinrichs A J (2009). Digestion and nitrogen utilization in dairy heifers limit-fed a low or high forage ration at four levels of nitrogen intake. *Journal of Dairy Science* **92**: 2078-2094