



Tarım Bilimleri Dergisi
Tar. Bil. Der.

Dergi web sayfası:
www.agri.ankara.edu.tr/dergi

Journal of Agricultural Sciences

Journal homepage:
www.agri.ankara.edu.tr/journal

Activity Guided Isolation of Nematicidal Constituents from the Roots of *Berberis brevissima* Jafri and *Berberis parkeriana* Schneid

Saqib ALI^a, Ishrat NAZ^b, Muhammad ALAMZEB^c, Mamoon URRASHID^d

^aDepartment of Chemistry, Faculty of Sciences, University of Kotli Azad Jammu and Kashmir, 11100, PAKISTAN

^bDepartment of Plant Pathology, The University of Agriculture, Peshawar, Khyber Pakhtunkhwa, 25130, PAKISTAN

^cDepartment of Chemistry, Faculty of Sciences, University of Kotli Azad Jammu and Kashmir, 11100, PAKISTAN

^dDepartment of Chemistry, Faculty of Arts and Basic Sciences, The Baluchistan University of IT, Engineering and Management Sciences (BUIITEMS), Takatu Campus, Quetta-87100, PAKISTAN

ARTICLE INFO

Research Article

Corresponding Author: Saqib ALI, E-mail: saqibali.chem@gmail.com, Tel: +92 (333) 984 48 42

Received: 03 November 2017, Received in Revised Form: 13 March 2018, Accepted: 17 April 2018

ABSTRACT

Biological screening of different parts of the selected *Berberis* species (*B. brevissima* Jafri and *B. parkeriana* Schneid) showed that methanolic root extract possessed significant efficacy against *Meloidogyne javanica* (a root knot nematode). From root methanolic extracts of selected *Berberis* species four isoquinoline alkaloids; jatrorrhizine, dehydrocheilanthifoline, berberine and berberrubine were isolated. Structures of the isolated compounds were determined by using EIMS, ¹H and ¹³C NMR, and other 2D spectroscopic techniques. Percentage juveniles mortality of *M. javanica* was determined at various concentrations (100, 200 and 300 µg mL⁻¹) using carbofuran as control. Berberine possessed the highest nematicidal activity (71.33%) followed by jatrorrhizine (59.50%). The *in vitro* results suggested that these compounds from *Berberis* species could be potential novel nematicides against *M. javanica*.

Keywords: Nematicidal activity; *Berberis brevissima*; *Berberis parkeriana*; *Meloidogyne javanica* and isoquinoline alkaloids

© Ankara Üniversitesi Ziraat Fakültesi

1. Introduction

All over the world, in the field of agriculture root knot nematodes (*Meloidogyne* species) are mainly responsible for huge economic losses (Liu et al 2011). For several decades the use of various chemical nematicides is an important tool to control root knot nematodes. But due to its negative impact on environment and after use resistances, have either reduced or totally banned its use, therefore these nematicides must be replaced with safe and

more effective chemical nematicides (Zuckerman & Esnard 1994). In the area of vegetables and fruits production, approximately 70 billion U.S. dollar crop damage is due to these root knot nematodes (*Meloidogyne* species) annually (Reynolds et al 2011). Amongst the all probable strategies for controlling these pests, the biocontrol agents obtained from plant or microorganisms could be used to lower non-target contact of harmful pesticides and to face resistance growth (Isman 2006; Tian et al 2007). Different types of plants, constituents and

metabolites have been screened for efficacy against various plant nematodes (Hong et al 2007; Thoden et al 2009; Ntalli et al 2010; Bai et al 2011).

Berberis (Berberidaceae) possesses more than 500 species and is the only genus of the family in the southern hemisphere (Bai et al 2011). The genus *Berberis* is full of isoquinoline alkaloids having high potential in the treatments of many ailments and insects control (Baird et al 1997; Wright et al 2000; Quevedo et al 2008). The hydro ethanolic extract of *Berberis* species (*B. aristata*, *B. asiatica*, *B. chitria* and *B. lyceum*) have shown very good antimicrobial efficacy against bacterial (eleven) and fungal (eight) strains (Küpeli et al 2002). In scurvy, angina, sore throat and dysentery its leaves decoction has been used as antiscorbutic. Berries of the genus could be used as a tonic and be used in the form of a dye (Singh et al 2007). Various Chinese folk remedies have reported use of different species of the genus *Berberis* (*B. aquifolium*, *B. aristata* and *B. vulgaris*) for inflammations and rheumatic problems (Li et al 1989; Ju et al 1990; Teh et al 1990; Kondo et al 1992; Saied & Begum 2004). *B. aristata* has shown a very high anticancer efficacy against colon cancer cell line (HT29) (Seow et al 1992). In the current study, we have isolated four isoquinoline alkaloids through activity guided fractionation of methanolic crude extract of roots of *Berberis* species (*B. brevissima* and *B. parkeriana*) and studied their efficacy against root knot nematode *M. javanica*.

2. Material and Methods

2.1. General

Silica gel 60 (0.063-0.200 mm) was used for Column Chromatography (CC) while silica gel 60 PF254 was used for preparative Thin Layer Chromatography (TLC). The melting points of isolated compounds were determined by the melting point apparatus (Bibby Scientific Limited, Stone Staffordshire ST15 0SA, UK). UV spectra were taken by Thermo Spectronic UNICAM UV 300. IR spectra were recorded using JAEKO FT/IR-4200/A. Spectral characterizations of the compounds were performed by using Bruker AVANCE 500 and 400 MHz

instruments. ^{13}C Nuclear Magnetic Resonance spectra (^{13}C NMR) were recorded at 100 MHz while ^1H Nuclear Magnetic Resonance (^1H NMR) spectra at 500 MHz and 400 MHz using deuterated chloroform and methanol as solvents. EIMS (JEOL MSRoute) was determined by using direct insertion probe.

2.2. Plant material

B. brevissima roots (2.5 kg) were collected from Tirah (Khyber Agency, Khyber Pakhtunkhwa, Pakistan) and *B. parkeriana* roots (2 kg) from Dir (Lower) (Khyber Pakhtunkhwa). The species were identified by Prof. Dr. Jandar SHAH (Ex. Voice Chancellor Benazir Bhutto University, Sherengal, Khyber Pakhtunkhwa, Pakistan). The voucher specimens (No. Bot/10710 and 8719) were deposited in herbarium (Department of Botany, University of Peshawar).

2.3. Extraction and chromatography

The plant material was soaked in 95% methanol for 7 days and the solvent was then evaporated at 40 °C (reduced pressure), using rotary evaporator. The residue obtained (*B. brevissima* root methanolic extract (BBR-MeOH= 225.7 g) and *B. parkeriana* root methanolic extract (BPR-MeOH= 186.5 g)) were dissolved in 6.5 L of 5% acidic water, filtered and left over night at room temperature. The yellow precipitate was filtered to obtain fraction A (*B. brevissima* root fraction A (BBR-FA= 86.3 g) and *B. parkeriana* root fraction A (BPR-FA= 77.5 g)). The filtrate was then extracted with CH_2Cl_2 (3x300 mL). The organic layer was separated and evaporated to afford fraction B (*B. brevissima* root fraction B (BBR-FB= 8.5 g) and *B. parkeriana* root fraction B (BPR-FB= 6.0 g)). The aqueous (acidic) layer was basified (pH 9-10) with concentrated aqueous NH_3 and then extracted with CHCl_3 (3x300 mL). Which was evaporated to afford fraction C (*B. brevissima* root fraction C (BBR-FC= 17.5 g) and *B. parkeriana* root fraction C (BPR-FC= 14.8 g)). The remaining aqueous layer was dried to obtain fraction D (*B. brevissima* root fraction D (BBR-FD= 53.5 g) and *B. parkeriana* root fraction D (BPR-FD= 45.8 g)).

2.3.1. Fraction A

On recrystallization of fraction A (BBR-FA and BPR-FA) almost pure berberine (3) (56 g) was obtained, mp 207-209 °C (lit 208-210 °C) (Suau et al 1998).

2.3.2. Fraction C

BBR-FC was further put to column chromatography (Silica gel, 180 g) and eluted with CHCl_3 -MeOH with increasing polarity. The fractions so obtained were further subjected to preparative TLC using CHCl_3 -MeOH- NH_3 (90:10:0.5 and 85:15:0.5) to give dehydrocheilanthifoline (2) (15.3 mg) and Jatrorrhizine (1) (21.6 mg) respectively. BPR-FC was also separated on CC (Silica gel, 150 g) eluted with CHCl_3 -MeOH (10:90) followed by preparative TLC using CHCl_3 -MeOH- NH_3 (10:90:0.5) to gave berberrubine (4) (7.5 mg).

2.4. Nematicidal assay

Pure culture of *M. javanica* was obtained from Department of Plant Pathology, The University of agriculture, Peshawar, Khyber Pakhtunkhwa and was maintained on tomato cultivar Rio-grande through single egg mass inoculation. For 60 days the tomato plants were grown inside the glass house. *M. Javanica* eggs were extracted (1% NaOCl solution) rinsed with on 25 μm aperture sieve (distilled water). Juveniles of second stage (J2s) were obtained from surface sterilized eggs placed in sterile water in a cavity block, which hatched after 3-4 days (Hussey & Barker 1973). Nematicidal assay of the various fractions and pure compounds were determined against *M. javanica* (J2s) using microwell assay (Naz et al 2012). All fractions and pure compound's stock solutions were prepared by dissolving them in 1% DMSO and was further diluted using distilled water. From the slandered solutions (300 $\mu\text{g mL}^{-1}$) final concentrations (100, 200 and 300 $\mu\text{g mL}^{-1}$) were prepared (Naz et al 2012). Second stage juveniles (100) were transferred to 24 microwell plate (Multiwell, TM 24, Becton Dickinson, USA) in final volume of 1 mL in various concentrations (100, 200 and 300 $\mu\text{g mL}^{-1}$) of the fractions and pure compounds, carbofuran (with

DMSO; 1% v/v) was used as a positive control. At room temperature (25 °C), the experiment was performed twice and repeated four times for each concentrations. After 24 hours of incubation, total number of active or inactive J2s were recorded. Finally to find out the mobility or mortality after 24 hours, the J2s was transferred to distilled water. The J2s were considered as dead if they did not move even after mechanical prodding (Choi et al 2007). Percentages of J2s mortality was calculated for each well and the results obtained were subjected to ANOVA (Analysis of Variance) and means were separated through Fisher's projected least significance difference (LSD) test at $P= 0.05$ using MSTAT-C software (Gomez & Gomez 1984).

3. Results and Discussion

3.1. Activity guided isolated bioactive compounds

Four isoquinoline alkaloids were isolated through activity guided fractionation of methanolic crude extracts and their structures were characterized by various spectroscopic techniques. The values were compared with the literature data and the structures are given in Figure 1.




Figure 1- Structures of isolated nematicidal bioactive compounds

3.2. Compound characterization

3.2.1. Jatrorrhizine

Brown crystalline compound (MeOH); melting point: 281-282 °C (lit 280-282 °C) (Hsieh et al 2004). Molecular formula $C_{20}H_{20}N^+O_4$, EIMS m/z: 338.1387 (M^+). UV λ_{max} nm (MeOH) 226.0, 265.0, 349.0, 435.5; IR ν_{max} cm^{-1} 3340.1, 2942.8, 1600.6. The melting point, EIMS, UV, IR, 1H and ^{13}C NMR data were in agreement with the literature (Hsieh et al 2004).

3.2.2. Dehydrocheilanthifoline

Brown amorphous compound (MeOH), melting point 269-270 °C. Molecular formula $C_{19}H_{16}N^+O_4$, deduced from the EIMS m/z 322.1074. UV λ_{max} nm (MeOH) 264.5, 359.0, 464.0; IR ν_{max} cm^{-1} 3361.8, 2924.5, 2358.5, 1601.6 (Santavy 1979).

3.2.3. Berberine

Yellow crystalline compound (MeOH), melting point 207-09 °C (lit 208-210 °C) (Suau et al 1998). Molecular formula $C_{20}H_{18}N^+O_4$, EIMS m/z: 336.1230 (M^+). UV λ_{max} nm (MeOH) 264.50, 349.0, 427.5; IR ν_{max} cm^{-1} 3047.9, 2925.5, 1596.8. The melting point, EIMS, UV, IR, 1H and ^{13}C NMR data were in agreement with the literature (Hsieh et al 2004).

3.2.4. Berberrubine

Brown amorphous compound (MeOH), melting point 257-260 °C (lit. 255-259 °C) (Liu et al 2010). Molecular formula $C_{19}H_{15}NO_4$, EIMS m/z: 321.1001 (M^+), 306.3 (M^+-15), 292.3, 278.3. The EIMS, 1H and ^{13}C NMR values were in close similarity to the reported one (Shamma & Rahimizadeh 1986).

3.2.4.1. In vitro nematocidal efficacy of crude extracts and various fractions obtained from the berberis species against J2s mortality of *M. javanica*

The data obtained (Table 1) revealed significant ($P \leq 0.05$) effect on mortality of J2s at various concentrations i.e. 100, 200 and 300 $\mu g mL^{-1}$. The mortality of J2s was increased with the increase in concentration, the highest concentration (300 $\mu g mL^{-1}$) was most effective (54.10%). *Berberis brevissima* roots methanolic crude extract (BBR-MeOH) showed 22.33% J2s mortality and *Berberis parkeriana* methanolic roots extract (BPR-MeOH) 31.11%. Amongst the different fractions BBR-FA showed the highest mortality of 62.22% followed by BPR-FA (57.22%) and BBR-FB (54.00%). The BBR-FB and BPR-FB exhibited approximately 50% activity of the standard (carbofuran) but the differences were non-significant (Table 1).

Table 1- In vitro effect of different concentrations of plant fractions on juvenile mortality of root knot nematode *Meloidogyne javanica*^a

Fractions	Concentrations			Mean
	100 $\mu g mL^{-1}$	200 $\mu g mL^{-1}$	300 $\mu g mL^{-1}$	
BBR-MeOH	18.33	23.00	25.67	22.33 g
BBR-FA	56.33	59.00	71.33	62.22 b
BBR-FB	41.67	48.67	51.66	47.33 d
BBR-FC	50.67	62.33	78.33	54.00 c
BBR-FD	39.00	43.33	45.67	42.67 e
BPR-MeOH	25.33	31.00	37.00	31.11 f
BPR-FA	35.67	40.67	44.00	40.11 e
BPR-FB	45.00	46.67	47.00	46.22 d
BPR-FC	54.00	58.00	59.00	57.22c
Carbofuran	89.67	89.67	96.67	88.22 a
Mean	44.27 c	49.07 b	54.10 a	

Data are means of five replicate per treatment using the combination of two experiments (Spring and fall, 2011); ^a, means followed by the same letters do not differ significantly ($P \leq 0.05$) according to Fisher's protected LSD test. (LSD value for fractions= 3.31, LSD value for concentration= 1.82, LSD value for interaction= 6.62)

3.2.4.2. *In vitro* nematicidal activity of isoquinoline alkaloids from the two berberis spp. against second stage juvenile mortality of *M. javanica*

In vitro nematicidal efficacy of the four isolated alkaloids at various concentrations (100, 200 and 300 $\mu\text{g mL}^{-1}$) and its interaction were determined ($P \leq 0.05$). Increase in mortality of J2s was linear ($R^2 = 0.98$) dose dependent (Figure 2). Second stage juvenile mortality was 76.67% at a concentration of 300 $\mu\text{g mL}^{-1}$. Figure 3 indicated significant effect of the pure compounds at various concentrations. Amongst the tested compounds berberine (3) exhibited highest potential (97.3%) of the standard carbofuran at a concentration of 300 $\mu\text{g mL}^{-1}$. Amongst the isolated compounds jatrorrhizine (1) ranked second with efficacy of 59.50% followed by berberrubine (4) with J2s mortality of 49.17% (Figure 3). In the four isolated isoquinoline alkaloids dehydrocheilanthifoline (2) was less effective, nevertheless, showed significant mortality at the tested concentrations. The interaction of isolated compounds were studied at various concentrations and were lightly significant (Figure 4). The data showed that the percentage J2s mortality increased as the concentration of tested compounds were increased (Figure 4).

Literature survey indicated the antibacterial activity of the alkaloids of *B. thunbergii* DC and *B. vulgaris* (L) (Villinski et al 2003). The stem bark of *B. asiatica* L. showed high antimicrobial activity than the standard (Bhandari et al 2000), while the fresh and dried, aqueous as well as methanolic extracts of *B. asiatica* showed good activity against G-positive and G-negative bacteria (Shahid et al 2009). Berberine was suggested to be the main antimicrobial component of the plant. Alkaloids was suggested to have microbiocidal properties (Ghoshal et al 1996) whereas berberine has been found effective against many trypanosomes (Freiburghaus et al 1996), plasmodia (Omulokoli et al 1997) and many invertebrate pests (Rattan 2010). It was suggested that mechanism of action of berberine could be attributed to its ability to intercalate with the DNA synthesis of parasites (Phillipson et al 1987).




Figure 2- Effect of three different concentrations of pure compounds of *Berberis* spp., on % J2s mortality of *Meloidogyne javanica*




Figure 3- *In vitro* nematicidal effect of different pure compounds of *Berberis* spp., against % J2s mortality of *Meloidogyne javanica* (Berb, Berberrubine; Jat, Jatrorrhizine; Ber, Berberine; Dehd, Dehydrocheilanthifoline)




Figure 4- *In vitro* interaction effect between pure compounds of *Berberis* spp., and three different concentrations on % J2s mortality of *Meloidogyne javanica* (Berb, Berberrubine; Jat, Jatrorrhizine; Ber, Berberine; Dehd, Dehydrocheilanthifoline)

4. Conclusions

In the present study we have found that fractions of methanolic extracts of the two *Berberis* species have high potential against the root knot nematodes. Secondary metabolites of plants could be used as defense (toxic), which hinder reproduction and other physiological and biological functions of pests and parasites. These biomolecules could be used for enhancing the effectiveness and specificity in future nematicides design with specific or multiple target sites. These studies suggest that methanolic crude extracts and especially the isoquinoline alkaloids could be used as potential novel nematicides against *M. javanica*. Further research encompassing the isolation and identification of more nematicidal isoquinoline alkaloids from *Berberis* spp. may be carried out and tested against root knot nematodes as well as other plant parasitic nematodes.

Acknowledgments

The authors thank to Higher Education Commission of Pakistan for financial support and Prof. Dr. Jandar SHAH for identification of the selected species.

References

- Bai C Q, Liu Z L & Liu Q Z (2011). Nematicidal constituents from the essential oil of *Chenopodium ambrosioides* aerial parts. *E-Journal of Chemistry* 8: 143-148
- Baird A W, Taylor C T & Brayden D J (1997). Non antibiotic anti diarrhoeal drugs: factor affecting oral bioavailability of berberine and loperamide in intestinal tissue. *Advanced Drug Delivery Review* 23(1-3): 111-120
- Bhandari N, Verma H C, Upadhyay C, Tripathi A & Pathi R P (2000). Mossbauer Spectroscopy of K/T boundary clays: Characteristics of iron bearing minerals; In: Catastrophic events and mass extinctions: Impacts and beyond, Houston: Lunar and Planetary Institute, pp. 12-13
- Choi W Y, Giraldez A J & Schier A F (2007). Target protectors reveal dampening and balancing of Nodal Agonist and Antagonist by miR-430. *Science* 318(5848): 271-274
- Freiburghaus S, Kaminsky R, Nkunya M H H & Brun R (1996). Evaluation of African medicinal plants for their *in-vitro* trypanocidal activity. *Journal of Ethnopharmacology* 55(1): 1-11
- Ghoshal S, Krishna B N & Lakshmi V (1996). Antiamoebic activity of *Piper longum* fruits against *Entamoeba histolytica* *in-vitro* and *in-vivo*. *Journal of Ethnopharmacology* 50(3): 167-170
- Gomez K A & Gomez A A (1984). Two factor experiment. In *Statistical procedures for agriculture research*, John Wiley and Sons: New York, USA, pp. 89-109
- Hong L J, Li G H, Zhou W, Wang W & Zhang K Q (2007). Screening and isolation of a nematocidal sesquiterpene from *Magnolia grandiflora* L. *Pest Management Sciences* 63(3): 301-305
- Hsieh T J, Chia Y C, Wu Y C & Chen C Y (2004). Chemical constituents from the stems of *Mahonia japonica*. *Journal of Chinese Chemical Society* 51(2): 443-446
- Hussey R S & Barker K R (1973). A comparison of methods of collecting inocula of *Meloidogyne* spp., including a new technique. *Plant Disease Report* 57: 1025-1028
- Isman M B (2006). Botanical insecticides, deterrents, and repellents in modern agriculture and an increasingly regulated world. *Annual Review of Entomology* 51: 45-66
- Ju H S, Li X J, Zhao B L, Han Z W & Xin W J (1990). The scavenging effect of berbarmineon active oxygen radicals in phorbol ester stimulated human polymorphonuclear leukocytes. *Biochemical Pharmacology* 39(11): 1673-1678
- Kondo Y, Imai Y, Hojo H, Hashimoto Y & Nozoe S (1992). Selective inhibition of T cell dependent immune responses by bisbenzylisoquinoline alkaloids *in vivo*. *International Journal of Immunopharmacology* 14(7): 1181-1186
- Küpeli E, Koşar M, Yeşilada E, Hüsni K & Başer C (2002). A comparative study on the antiinflammatory, antinociceptive and antipyretic effects of isoquinoline alkaloids from the roots of Turkish *Berberis* species. *Life Sciences* 72(6): 645-657
- Li S Y, Ling L H, Teh B S, Seow W K & Thong Y H (1989). Antiinflammatory and immunosuppressive properties of the bisbenzylisoquinolines: *In vitro* comparisons of tetrandine and berberine. *International Journal of Immunopharmacology* 11(4): 395-401

- Liu Y T, Hao H P, Xie H G, Lai L, Wang Q, Liu C X & Wang G J (2010). Extensive intestinal first pass elimination and predominant hepatic distribution of berberine explain its low plasma levels in rats. *Drug Metabolism and Disposition* 38(10): 1779-1784
- Liu J H, Wang L, Qiu J Y, Jiang L I, Yan J Y, Liu T, Liu W C & Duan Y X (2011). Nematocidal activity of *Gymnoascus reesii* against *Meloidogyne incognita*. *African Journal of Microbiology Research* 5(18): 2715-2719
- Naz I, Rius J E, Saifullah, Blok V, Khan S R, Ali S & Ali S (2012). *In vitro* and in planta nematocidal activity of *Fumaria parviflora* (Fumariaceae) against Southern Root knot nematode *Meloidogyne incognita*. *Plant Pathology* 62: 943-952
- Ntalli N G, Ferrari F, Giannakou I & Spiroudi M (2010). Phytochemistry and nematocidal activity of the essential oils from 8 greek lamiaceae aromatic plants and 13 terpene components. *Journal of Agriculture and Food Chemistry* 58(13): 7856-7863
- Omulokoli E, Khan B & Chhabra S C (1997). Antiplasmodial activity of four Kenyan medicinal plants. *Journal of Ethnopharmacology* 56(2): 133-137
- Phillipson O T, Kilpatrick I C & Jones M W (1987). Dopaminergic innervation of the primary visual cortex in the rat, and some correlations with human cortex. *Brain Research Bulletin* 18: 621-633
- Quevedo R, Valderrama K, Murillo M B, Laverde M & Fajardo V (2008). A new bisbenzyltetrahydroisoquinoline alkaloid from *Berberis tabiensis* (Berberidaceae). *Biochemical Systematics and Ecology* 36(10): 812-814
- Rattan R S (2010). Mechanism of action of insecticidal secondary metabolites of plant origin. *Crop Protection* 29(9): 913-920
- Reynolds A M, Dutta T K, Curtis R S C, Power S J, Gaur H S & Kerry B R (2011). Chemotaxis can take plant parasitic nematodes to the source of a chemo attractant via the shortest possible routes. *Journal of the Royal Society Interface* 8: 568-577
- Saied S & Begum S (2004). Phytochemical studies of *Berberis vulgaris*. *Chemistry of Natural Compounds* 40(2): 137-140
- Santavy F (1979). Berbine (dibenzo[a,g]quinolizidine) (protoberberine, pseudoprotoberberine, corydaline and corytenchirine types) group. In *The Alkaloids*, Manske R H F Eds.; Academic Press: New York, USA, 17, pp. 439-461
- Seow W K, Ferrante A, Summors A & Thong Y H (1992). Comparative effects of tetrandrine and berbamine on production of the inflammatory cytokines interleukin-1 and tumour necrosis factor. *Life Sciences* 50(8): PL53-PL58
- Shahid M, Rahim T, Shahzad A, Tajuddin, Latif A, Fatma T, Rashid M, Raza A & Mustafa S (2009). Ethnobotanical studies on *Berberis aristata* DC. root extracts. *African Journal of Biotechnology* 8(4): 556-563
- Shamma M & Rahimizadeh M (1986). The identity of chileninone with berberrubine. The problem of true natural products vs. artifacts of isolation. *Journal of Natural Products* 49(3): 398-405
- Singh M, Srivastava S & Rawat A K S (2007). Antimicrobial activities of Indian *Berberis* species. *Fitoterapia* 78(7): 574-576
- Suau R, Rico R, Romero J M L, Najera F & Cuevas A (1998). Isoquinoline alkaloids from *Berberis vulgaris* Subsp. *australis*. *Phytochemistry* 49(8): 2545-2549
- Teh B S, Seow W K, Li S Y & Thong Y H (1990). Inhibition of prostaglandin and leukotriene generation by the plant alkaloids tetrandrine and berbamine. *International Journal of Immunopharmacology* 12(3): 321-326
- Thoden T C, Boppre M & Hallmann J (2009). Effects of pyrrolizidine alkaloids on the presence of plant parasitic and free living nematodes. *Pest Management Sciences* 65(7): 823-830
- Tian B, Yang J & Zhang K Q (2007). Bacteria used in the biological control of plant parasitic nematodes: Populations, mechanisms of action, and future prospects. *FEMS Microbiology Ecology* 61(2): 197-213
- Villinski J, Dumas E, Chai H B, Pezzuto J, Angerhofer C & Gafner S (2003). Antibacterial activity and alkaloid content of *Berberis thunbergii*, *Berberis vulgaris* and *Hydrastis Canadensis*. *Pharmaceutical Biology* 41(8): 551-557
- Wright C W, Marshall S J, Russell P F, Anderson M M, Phillipson J D, Kirby G C, Warharst D C & Schiff J P L (2000). In vitro antiplasmodial, antiamebic and cytotoxic activities of some monomeric isoquinoline alkaloids. *Journal of Natural Products* 63(12): 1638-1640
- Zuckerman B M & Esnard J (1994). Biological control of plant nematodes current status and hypothesis. *Japanese Journal of Nematology* 24: 1-13